

AD695177

FSTC-HT-23-443-69

# U.S. ARMY FOREIGN SCIENCE AND TECHNOLOGY CENTER




THE STUDY OF THE CONCENTRATIONS OF LONG-DURATION  
RADIOACTIVE AEROSOLS OF ATMOSPHERIC  
AIR IN ITS RELATION TO THE AMOUNT OF DUST

COUNTRY: USSR

## TECHNICAL TRANSLATION

This document has been approved  
for public release and sale;  
its distribution is unlimited.  
It may be released to the  
Clearinghouse, Department of  
Commerce, for sale to the  
general public.



TECHNICAL TRANSLATION

FSTC-HT-23- 443-69

THE STUDY OF THE CONCENTRATIONS OF LONG-DURATION  
RADIOACTIVE AEROSOLS OF ATMOSPHERIC  
AIR IN ITS RELATION TO THE AMOUNT OF DUST

by

M. V. KOZLOVA, Candidate of Technical Science,  
I. P. KORENKOV, Candidate of Medical Science,  
YU. V. NOVIKOV and D. D. GRANOVSKAYA

The Moscow Municipal Epidemic Hospital and The  
Moscow Scientific Research Institute of Hygiene  
called F. F. Erisman  
USSR

Translated for FSTC by Techtran Corporation

This translation is an unedited rendition of the original foreign text. Statements or theories advocated or implied are those of the source and do not reflect the position or opinion of the US Army Foreign Science and Technology Center. This translation is published with a minimum of copy editing and graphics preparation in order to expedite the dissemination of information. Requests for additional copies of this document should be addressed to the Defense Documentation Center, Cameron Station, Alexandria, Virginia, ATTN: OSR-2.

THE STUDY OF THE CONCENTRATIONS OF LONG-DURATION  
RADIOACTIVE AEROSOL OF ATMOSPHERIC  
AIR IN ITS RELATION TO THE AMOUNT OF DUST

Radioactive substances are found in the air in the form of aerosols. The diameter of the latter particles is equal to tenths and hundreds of a micron. These small particles of the products of division, which are carried great distances by air currents, settle out on larger ones which are always present in the surface layer of the atmosphere of air, converting them in this way to radioactive particles (N. A. Susorova; Eisenbud and Harley).

We were not able to find data in technical literature on the study of the correlation between the concentrations of long-duration aerosols in the surface layer of the atmosphere and the amount of dust in it. In addition, this question has a certain hygienic significance. It is well known that an increase in the concentration of atmospheric dust containing stable isotopes can lead to a decrease in the specific activity of dust hovering in the air. In the given case, stable atmospheric dust appears in the roll of a factor which reduces the amount of radioactive substances in the atmosphere of air. Since it is necessary to take this condition into consideration in the hygienic evaluation of the radioactive contamination of the atmospheric air, we carried out as an experiment, a determination of the concentration of long-duration radioactive substances and dust of the atmospheric air in Moscow.

We took samples with the help of a special aspiration device UAS-1 (V. P. Rublevskiy) equipped with a FPP-15 filter. The area of the filter was 5500 cm<sup>2</sup>. The volume of filtered air on the average came to 1200 m<sup>3</sup>. During 1965 seventy-eight samples of atmospheric air were taken for tests of radioactivity and the content of stable dust at an altitude of 15 meters over soil covered with asphalt.

The concentration of dust in one cubic meter of air was determined by preliminary weighing of the filter after the action of the aspirator was turned off. Weighing was accomplished after first leaving the filter in a desiccator for two days. In addition, the filter was heated in the muffle furnace at 400°C. With the aim of determining the concentration of long-duration  $\beta$ -active aerosols, we took a weight sample of 10 mg of an obtained quantity of ash and applied it to the standard aluminum footboard; then on the DP-100 device with a BFL T-25 counter, we determined the total activity of the samples being investigated.

The specific  $\beta$ -activity of the long-duration aerosols (A) in curie for one liter was determined according to the following formula:

$$A = \frac{(N - N_{\phi}) \cdot p_2}{2.2 \cdot 10^{10} \cdot V \cdot k \cdot p_1}$$

where N is the rate of calculation (per minute) from the samples being investigated,  $N_{\phi}$  is the phon of the device (per minute), v is the volume of the air drawn in (in liters); k is the effectiveness of the counting mechanism determined with the help of  $K^{40}$  (N. A. Susorova),  $p_1$  is the correction for self-absorption in the preparation,  $p_2$  is the correction for brake-through of the aerosols by means of the FPP-15 (10%).

The results of the investigation of the concentrations of  $\beta$ -active long-duration aerosols and dust of the atmospheric air are shown in the figure. From this figure it is clear that the amount of contamination of the atmospheric air by  $\beta$ -active long-duration aerosols in 1965 was at a level of  $\sim 1.9 \cdot 10^{-17}$  curie/per liter. The concentration of dust in the surface layer of the atmosphere was in the range of 0.03 - 0.7 mg/m<sup>3</sup>. Judging from the drawing, a direct relationship of the concentrations of the  $\beta$ -active aerosols to the amount of dust in the air was noted at different

months. For example, on the 13th of January, 12th of April, the 23rd and 28th of July, and on other days when there was an increase in the amount of dust in the atmospheric air, the concentrations of radioactive aerosols in it increased. In other cases (on 31 March, 19 April, 5 May etc.) such a relationship was not observed.

Taking these data into consideration, we set out to determine the relationship between the concentrations of radioactive substances and the amount of dust in atmospheric air by using the method of correlation analysis (A. M. Dlin). With this aim, we used the correlation grid for results of measurement and constructed the empirical line of regression which was expressed by the equation:

$$y_x = f(x),$$

where  $y$  is the average concentration of radioactive substances in atmospheric air,  $x$  is the amount of dust in the air.

The theoretical line of regression has the form:

$$y = ax + b$$

where  $y$  is the concentration of the radioactive aerosols (in curie/liter),  $x$  is the amount of dust in the air (in  $\text{mg}/\text{m}^3$ ),  $a$  and  $b$  are the coefficients for the equation of regression;  $a$  is the amount of dust in the air (in  $\text{mg}/\text{m}^3$ ) at a concentration of radioactive substances (in curie/liter),  $b$  is the concentration of radioactive substances in the air in the absence of dust.

In our investigations, the analytic expression for the theoretical line of regression had the form:

$$y = 6,6x + 3,5,$$

where  $a = 6.6$ ,  $b = 3.5$ .




Fig. 1. Concentration of long-duration  $\beta$ -active aerosols (1) and hovering dust (2) in the atmospheric air of Moscow in 1965.

Key: a. Concentration of radioactive substances in the atmospheric air (curie/liter); b. Jan., Feb. March, April, May, June, July Aug., Sept., Oct., Nov., Dec; c. Amount of dust of the atmospheric air (in mg/m<sup>3</sup>).

The coefficient of correlation was determined according to the formula:

$$r = a \frac{\sigma_x}{\sigma_y},$$

$$\text{where: } \sigma_x = h_x \cdot \sqrt{\frac{\sum m_x \cdot x'^2}{\sum m_x} - \left(\frac{\sum m_x \cdot x'}{\sum m_x}\right)^2},$$

$$\sigma_y = h_y \cdot \sqrt{\frac{\sum m_y \cdot y'^2}{\sum m_y} - \left(\frac{\sum m_y \cdot y'}{\sum m_y}\right)^2}.$$

$\sigma_x$  and  $\sigma_y$  is the dispersion of the values  $x$  and  $y$ ,  $h_x$  is the step of the integrals according to  $x$ ,  $h_y$  is the step of the integrals according to  $y$ ,  $m_x$  and  $m_y$  are partial sums.

Introducing the data from the correlation tables, we find the coefficient of correlation. In our case, it was equal to  $+0.4 \pm 0.08$ .

Thus, a correlation relationship was found between the amount of dust of atmospheric air and the concentrations of long-duration aerosols. This mutual relationship expressed by the coefficient of correlation with a plus sign, has a proportional character. In spite of the fact that the coefficient of correlation obtained by us was not very close to unity, the error in determining it amounts to  $\sigma_r = \pm 0.05$ , which points to an existing mutual relationship.

#### BIBLIOGRAPHY

1. Dlin, A. M., *Matematicheskaya Statistika v Tekhnike*, Moscow, 1958.
2. Rublevskiy, V. P., *Gigenita and Sanatgagsiya*, No. 9, p. 60, 1964.
3. Susorova, N. A., *Radioaktivnye Zagryazneniya Vneshney Sredy*, Moscow, p. 9, 1962.
4. Eisenbud, M. and Harley, J. H., *Science*, vol. 124, p. 251, 1956.

UNCLASSIFIED

Security Classification

## DOCUMENT CONTROL DATA - R &amp; D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Foreign Science and Technology Center US Army Materiel Command Department of the Army		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
3. REPORT TITLE  THE STUDY OF THE CONCENTRATIONS OF LONG-DURATION RADIOACTIVE AEROSOLS OF ATMOSPHERIC AIR IN ITS RELATION TO THE AMOUNT OF DUST		2b. GROUP	
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Translation			
5. AUTHOR(S) (First name, middle initial, last name)  M. V. Kozlova, Candidate of Technical Science; I. P. Korenkov, Candidate of Medical Science; Yu. V. Novikov and D. D. Granovskaya			
6. REPORT DATE 18 September 1969	7a. TOTAL NO. OF PAGES 5	7b. NO. OF REFS N/A	
8a. CONTRACT OR GRANT NO.  a. PROJECT NO.  02RD500 2301  d. USAECOM (WSMR)		8b. ORIGINATOR'S REPORT NUMBER(S)  FSTC-HT-23-443-69  9b. OTHER REPORT NUM(S) (Any other numbers that may be assigned this report)  ACSI Control Number (None)	
10. DISTRIBUTION STATEMENT  This document has been approved for public release and sale; its distribution is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY US Army Foreign Science and Technology Center	
13. ABSTRACT  This article presents materials on the study on the mutual relationship between concentrations of long-duration aerosols and the amount of dust in atmospheric air. It was found that this mutual relationship has a proportional -haracter.			

DD FORM 1473-1-73

UNCLASSIFIED  
Security Classification



**UNCLASSIFIED**  
Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Radioactive aerosols Atmospheric air						

**UNCLASSIFIED**  
Security Classification